



Clearly different.
Clearly better.

Women's Health | Infectious Disease | Cardiovascular | Drugs of Abuse | GI Disorders

Clearview® HIV 1/2 STAT-PAK®

Clearview HIV 1/2 STAT-PAK

A rapid, qualitative screening test for the detection of antibodies to HIV 1 and HIV 2 in human whole blood, serum, and plasma.



CLIA waived

Simple, two-step procedure for whole blood.

Clearly different

Provides highly sensitive and specific tests for the detection of human immunodeficiency virus (HIV) in whole blood.

- **Rapid Reporting:** Results in 15 minutes.
- **Reliability:** Built-in control confirms sample addition and proper test performance.
- **Flexibility:** Use with multiple sample types – whole blood (fingerstick or venipuncture), serum or plasma.

Clearly better

- **Reliable results** – 99.7% sensitive, 99.9% specific
- **Detects HIV 1 and HIV 2** for a complete picture in one test.
- **15 minute results** mean STAT decision making, quickly and confidently while patient is present.
- **Minimal sample size of 5 µL** reduces sample handling and exposure risk, less invasive for patient.
- **Multiple specimen types** to fit your lab and practice patterns.
- **Long 24-month shelf life** allows your offices to purchase in volume.



**Clearview HIV 1/2
STAT-PAK
20 Test Kit**
Order No. 92110


Whole Blood - CLIA Waived
Serum, Plasma -
Non-waived

**Clearview HIV 1/2
Control Kit**
Order No. 92112

CDC Guidelines – Then & Now


Then	Now
Targeted testing based on risk assesment	HIV screening is recommended for patients ages 13 to 64 in all healthcare settings
Specific, signed consent for HIV testing is required	Opt-out required
HIV screening should be included in the routine prenatal screening tests for all pregnant women	HIV screening should be included in the routine prenatal screening tests for all pregnant women
Routinely recommended HIV testing for all patients seeking STD treatment	Persons at high risk for HIV infection should be screened for HIV at least annually
Pre-test counseling was required	Prevention counseling should not be required with HIV diagnostic testing, or as part of HIV screening programs in health care settings.

Awareness and Serostatus Among People with HIV and Estimates of Transmission ¹




- People living with HIV/AIDS: 1,039,000 - 1,185,000
- New sexual infections each year: ~32,000

HIV/AIDS Diagnoses among Adults and Adolescents, by Transmission Category 33 States, 2001 – 2004 ²



Clearview HIV 1/2 STAT-PAK Procedure



* Men having sex with men ** Intravenous drug user

1. Marks, et al. AIDS 2006; 20:1447-50
2. MMWR, Nov 18, 2005

All trademarks referenced are trademarks of their respective owners. All rights reserved.

visit
www.invernessmedicalpd.com
for more information